



2. Andrejewski Vorlesung/ SFB Seminartag

ZEIT:

10.1.2006, 14:00 Uhr - 19:00 Uhr

ORT:

im AEI in Golm, Max-Planck-Campus
Am Mühlenberg 1, 14476 Golm
Central Building, room number Z-050

PROGRAMM:

14:00 **Prof. Dr. Albrecht Klemm, University of Wisconsin,
Madison**

Mirror symmetry and the topological A- and B-model

Mirror symmetry on CY manifolds exchanges the symplectic structure on M , actually a complexified Kähler structure, with the complex structure on a mirror dual CY manifold W . The deformation theory of each of these structures can be described by a topological string theory called the topological A- and the B-model respectively. These models are cohomological theories defined by nilpotent operators Q_A and Q_B . We will show that Q_A exists on every symplectic manifold while Q_B exists only on CY manifolds and certain generalizations thereof. The latter fact is related to the Tian & Todorov theorem on the unobstructedness of complex structure deformations on CY spaces and generalizations by Hitchin. We will then discuss properties of cohomological theories notably the descend- and topological recursion relations. The solution of the topological B-model using these recursion relations and some classical methods of complex structure deformation theory on W is worked out and related by mirror symmetry to the Gromov-Witten theory captured by the topological A-model on M .

16:00 **Prof. Dr. Jens Hoppe (Royal Institute of Technology,
Stockholm), guest of the AEI**

Kontakt:

Humboldt-Universität zu Berlin . Institut für Mathematik
SFB 647 . Unter den Linden 6 . 10099 Berlin
Tel. +49 30 2093 1804 . Fax. +49 30 2093 2727
sfb647@math.hu-berlin.de

www.raumzeitmaterie.de

Aspects of Membrane Dynamics

17:30

Dr. Simon Chiossi (HU)

G2 structures on solvmanifolds

Conformally G2 manifolds are Riemannian manifolds with a G2 structure whose metric can be modified to a holonomy structure by a conformal change. There is an interesting construction of homogeneous conformally G2 structures on solvmanifolds built from underlying SU(3) structures. I will show how the corresponding non-homogeneous G2 metrics can be obtained also by evolving the SU(3) structure in time. (Possible reference: arXive math.DG/0510087)

Kontakt:

Humboldt-Universität zu Berlin . Institut für Mathematik
SFB 647 . Unter den Linden 6 . 10099 Berlin
Tel. +49 30 2093 1804 . Fax. +49 30 2093 2727
sfb647@math.hu-berlin.de

www.raumzeitmaterie.de